

Oil-in-Water Sunscreen Lotion SPF 30

Formula Number: JZ13-95

The formulation was subjected to *in vitro* analysis and delivered SPF ~35, PFA (PPD) ~34 performance with critical wavelength 381 nm.

| Phase | INCI Name [INN/USAN] (Other Information) | Trade Name | % Wt | Function |
|-------|---|--|-----------|--|
| A | Ethylhexyl Salicylate [Octisalate] | | 4.90 | UV Filter/Sunscreen Agent/Carrier System |
| | Ethylhexyl Methoxycrylene | SolaStay [®] S1 | 3.80 | Photostabilizer |
| | Butyloctyl Salicylate | HallBrite [®] BHB | 6.80 | Emollient/Sunscreen Solvent |
| | Glycine Soja (Soybean) Oil, Polyglyceryl-3 Diisostearate, Oryza Sativa (Rice) Germ Extract, Oryza Sativa (Rice) Extract | BLUE Oléoactif® | 1.00 | Active Ingredient |
| | Diisobutyl Adipate | Halltress [®] DIBA Special | 6.00 | Emollient/Sunscreen Solvent |
| В | Diethylamino Hydroxybenzoyl Hexyl Benzoate | | 4.00 | UV Filter/Sunscreen Agent |
| | Bis-Ethylhexyloxyphenol Methoxyphenyl Triazine | | 3.00 | UV Filter/Sunscreen Agent |
| | Butyl Methoxydibenzoylmethane [Avobenzone] | | 4.50 | UV Filter/Sunscreen Agent |
| с | VP/Eicosene Copolymer | | 2.00 | Film Former |
| | Cetyl Palmitate, Sorbitan Palmitate, Sorbitan Olivate | Oliwax [®] LC | 1.50 | Stabilizing Agent/Liquid Crystal Promoter |
| | Glyceryl Stearate | HallStar [®] GMS Pure | 1.00 | Emulsifier |
| | Cetearyl Olivate, Sorbitan Olivate | Olivem [®] 1000 | 2.00 | Self-Emulsifier/Liquid Crystal Maker |
| D | Water (Aqua) | | Up to 100 | Volatile Carrier |
| | Disodium EDTA | | 0.05 | Chelating Agent |
| Е | Glycerin | | 3.00 | Humectant |
| | Xanthan Gum (pharma grade) | | 0.20 | Thickener (Aqueous) |
| F | Potassium Cetyl Phosphate | | 1.40 | Emulsifier |
| G | Preservative | | a.n. | Preservative |
| | Dimethicone (5 cSt) | | 3.00 | Emollient |
| | Silica (spherical, 12 µm) | | 2.00 | Deoiling Agent |
| | Citric Acid | | a.n. | pH Adjuster |

PREPARATIVE PROCEDURE

- 1. To a secondary vessel add phase A and start mixing. Add phase B and heat to 80°C and then add phase C.
- 2. To the main vessel add phase D, start mixing and add premixture of phase E.
- 3. Heat phase DE to 75 80°C, and add phase F with mixing.
- 4. When both phases are homogeneous and at their designated temperatures, slowly add phase ABC to phase DEF with prop mixing.
- 5. After 5 minutes start cooling batch and when batch is < 50°C add phase E one by one with mixing
- 6. Continue mixing until batch is at room temperature, replacing any water lost to evaporation and mixing until smooth.
- 7. Adjust pH with 10%Citric acid to 5.0 5.0

PRODUCT CHARACTERISTICS (25°C)

Appearance (25°C): light yellow lotion

Viscosity (20 rpm, Brk, RV DV-II, T-D, after 24h at room temperature; mPa·s) 10000 - 16000